



2019
State of the Platte





Map depicting the Platte River system, including the Program Associated Habitat Reach on the central Platte River.



Program habitat complexes in the Associated Habitat Reach (AHR).

2019 State of the Platte

Adaptive Management Plan (AMP)
2019 “Big Question” Assessments*
February 20, 2020

*First Increment summary updated with 2007-2019 data.

Prepared by the Executive Director’s Office of the
Platte River Recovery Implementation Program
4111 4th Avenue, Suite 6
Kearney, NE 68845



PLATTE RIVER
RECOVERY IMPLEMENTATION PROGRAM



TABLE OF CONTENTS

5	Introduction
6	Summary of Key Learning from First Increment
9	Science Panel Comments on the 2019 State of the Platte
13	2019 Big Questions Table
14	2019 Big Question Assessments
34	Summary of Additional Learning
36	Appendix A - 2019 Priority Hypotheses Table
50	Appendix B - PRRIP Peer Reviewed Publications & Reports
49	Appendix C - PRRIP Program - Land & Water Objectives
51	Endnotes

INTRODUCTION

The Platte River Recovery Implementation Program’s (“Program” or “PRRIP”) Executive Director’s Office (EDO) developed this document for the Governance Committee (GC). It is intended to serve as a synthesis of Program monitoring data, research, analysis, and associated retrospective analyses to provide important information to the GC regarding key scientific and technical uncertainties. These uncertainties form the core structure of the Program’s Adaptive Management Plan (AMP) and are directly related to decisions regarding implementation of management actions, assessment of target species’ response to those management actions, how best the Program can spend its resources (money, land, water, etc.), and ultimately the success or failure of the Program.

A quick reference assessment for each of ten Big Questions is provided in Table 1, followed by an assessment write-up for each Big Question. Each assessment includes information noting any updates or changes from previous State of the Platte reports. This document contains endnotes to identify key documents or data sets that are important to read and understand when reviewing this report. Those endnotes include hyperlinks to information available in the Public Library section of the Program’s web site.

KEY OBSERVATIONS AND PROGRESS

The 2019 State of the Platte Report includes assessments incorporating Program data from years 2007-2018. Eight of ten Big Questions are answered conclusively, one (#3) is trending in a direction that will affirm or reject important hypotheses, and one question (#9) will be addressed by the GC during the AMP update process to be implemented during the First Increment Extension (2020-2032).

TERN AND PLOVER HABITAT AND USE

Implementation of short-duration high flow (SDHF) releases as currently envisioned will not create and/or maintain in-channel islands suitable for use and productivity for terns and plovers. In 2015 and 2016 the Program completed a Structured Decision Making (SDM) process to determine how to proceed with creating tern and plover habitat within the Associated Habitat Reach (AHR). Based on the SDM process, the GC completed the final “Adjust” stage of adaptive management and decided to maintain 10 acres of on-channel moving complex approach (MCA) islands and to create an additional 60 acres of off-channel nesting habitat.

WHOOPING CRANE HABITAT AND USE

Based on the findings of habitat selection analyses and related synthesis, the Program should continue management actions that provide unobstructed channel widths (UOCWs) for whooping cranes that are ≥650 ft and unforested corridor widths that are ≥1,100 ft. Implementation of SDHF releases as currently envisioned will not create

and/or maintain suitably-wide UOCW for whooping cranes. Mechanical methods such as disking and herbicide application at Program habitat complexes will create and maintain suitably wide UOCWs, though these management actions do not have the system-scale beneficial effects of natural flow events.

FORAGE AND TERN/PLOVER PRODUCTIVITY

Data analyses and syntheses indicate tern productivity is insensitive to flow and there is no evidence the forage base along the central Platte River limits tern and plover productivity. These same analyses and syntheses do not support Program summer flow releases to maintain the current 800 cubic feet per second (cfs) target for a diverse forage base for terns.

GLOSSARY OF ACRONYMS

ACRONYM	DEFINITION
AFY	Acre-feet per Year
AHR	Associated Habitat Reach
AMP	Adaptive Management Plan
AMWG	Adaptive Management Working Group
BQ	Big Question
cfs	Cubic Feet per Second
EDO	Executive Director’s Office
FSM	Flow-Sediment-Mechanical
GC	Governance Committee
ISAC	Independent Science Advisory Committee
LTPP	Least Tern and Piping Plover
MCM	Mechanical Creation and Maintenance
NF	Nearest Forest
PRRIP or Program	Platte River Recovery Implementation Program
SDHF	Short-Duration High Flow
SoPR	State of the Platte Report
TAC	Technical Advisory Committee
USFWS	United States Fish and Wildlife Service
UOCW	Unobstructed Channel Width
WAP	Water Action Plan
WC	Whooping Crane

Summary of Key Learning from AMP Implementation During the First Increment

As the Program begins the Extension in 2020, a review of First Increment learning relative to key AM components is in order to ensure Program AM activities during the Extension are focused on scientific and management uncertainties that are most relevant to the GC for decision-making. One key component is a set of **Management Objectives** that quantitatively describes tangible outcomes the Program is trying to achieve through implementation of the AMP. Performance against these management objectives is used to gauge the ability of the Program's management actions to provide benefits to the target species and achieve the Program Goal and Objectives. Another key component are **Management Strategies** which are logical packages of management actions that comprise a range of possible Program implementation learning activities on the ground. Summarizing First Increment learning through the lenses of these key components also reveals important **Remaining Uncertainties** that can be addressed through the implementation of AM during the Extension.

Management Objective #1 – Terns and Plovers

Objective – Improve production of the interior least tern and piping plover from the central Platte River. Proxy indicators include:

- a) Increase number of fledged tern and plover chicks
 - i) Increase nesting pairs (indicator is nesting pairs)
 - ii) Increase fledge ratios (indicator is chicks successfully produced per unit adult, nest or pair) and reduce chick mortality from causes such as flooding, predation, weather, inadequate forage.
- b) Reduce adult mortality
 - i) Reduce predation (indicator is nesting pairs)

First Increment Summary of Learning:

- Using the proxy indicators of fledge ratios and nesting pairs as specified in the AMP, the Program **met** Management Objective #1 during the First Increment.
- Least tern and piping plover populations in the AHR have increased significantly (Pearson's Rank Correlation test $\alpha \leq 0.05$) and proportionately to increases in habitat availability due to Program off-channel habitat creation efforts. Productivity on off-channel habitats has been sufficient to maintain a stable to growing subpopulation.
- Based upon available data, least tern and piping plover productivity is insensitive to river flow. Periods of low flow have not reduced productivity due to a limitation in forage availability.
- The Program agreed to acquire/develop and manage 60 more acres of off-channel tern and plover nesting habitat and 10 acres of MCA habitat to meet the Service's requirement of maintaining stable or growing tern and plover populations within the AHR.
- **Remaining Uncertainties** – need for and mechanics of additional predator control related to tern and plover productivity.

Management Objective #2 – Whooping Cranes

Objective – Contribute to the survival of whooping cranes during migration.

Proxy indicators include:

- a) Increase availability of whooping crane migration habitat along the central Platte River (indicators are the area of suitable roosting habitat, area of suitable foraging habitat, proportion of population, crane use days, etc.).

First Increment Summary of Learning:

- Using the proxy indicators of area of suitable roosting habitat, area of suitable foraging habitat, proportion of population, and crane use days as specified in the AMP, the Program **met** Management Objective #2 during the First Increment.
- Whooping crane use of the AHR has increased significantly (Pearson’s Rank Correlation test $\alpha \leq 0.05$) and proportionally to increases in habitat suitability during the spring migration season that are in part due to Program management actions.
- Whooping crane use of the AHR increased significantly (Pearson’s Rank Correlation test $\alpha \leq 0.05$) during the spring migration season while wet meadow use remained stable and low.
- **Remaining Uncertainties** – mechanics of flow releases (spring and fall migration flows, summer vegetation germination suppression flows) to ensure Program continues to meet the management objective.

Management Objective #3 – Pallid Sturgeon

Objective – Avoid adverse impacts from Program actions on pallid sturgeon populations.

- a) Indicators have not been identified as more research is needed to determine what potential indicators the Program may affect.

First Increment Summary of Learning:

- No proxy indicators for this management objective have been specified and therefore it is **unknown** if the Program met Management Objective #3 during the First Increment and learning related to pallid sturgeon and the Program has been limited.
- Translation of Program flow management actions from the central Platte to the lower Platte is difficult to detect and thus difficult to relate to effects on habitat and species response.
- **Remaining Uncertainties** – substantial uncertainty relating to the life history of pallid sturgeon in the lower Platte River (use, productivity, recruitment) limits the the ability of the Program to develop a clear set of testable hypotheses, management actions, monitoring protocols, and a plan for data analysis and synthesis.

Management Strategy #1 – Flow-Sediment-Mechanical (FSM)

Over-arching Hypothesis – combination of flow management, sediment management, and land management implemented concurrently will generate detectable changes in the channel morphology of the Platte River, and habitats for whooping crane, least tern, piping plover, pallid sturgeon, and other species of concern.

First Increment Summary of Learning:

- Attempts to implement the FSM management strategy have **generally produced poor results**.
- SDHF (5,000-8,000 cfs for three (3) days at Overton, NE) will not create or maintain suitable least tern and piping plover nesting habitat or whooping crane roosting habitat.
- Flow consolidation is not feasible due to legal and permitting constraints.
- A sediment deficit exists in the south channel downstream of the J-2 Return. Five to seven years of full-scale sediment augmentation are necessary to assess efficiency and effectiveness in preventing downstream migration of incision and narrowing.¹
- First Increment learning occurred largely through natural flow events as the Program was unable to implement a true SDHF and was not able to conduct flow consolidation actions.
- **Remaining uncertainties** – effectiveness of summer vegetation germination suppression flow and spring/fall WC migration flows in maintaining channel width.

Management Strategy #2 – Mechanical Creation & Maintenance (MCM)

Over-arching Hypothesis – combination of sandpit management, mechanical actions in the channel, and inundated wetlands and upland areas implemented concurrently will generate detectable changes in the channel morphology of the Platte River, and habitats for whooping crane, least tern, piping plover, pallid sturgeon, and other species of concern.

First Increment Summary of Learning:

- Implementation of the MCM management strategy has **produced mixed results**.
- Suitable on-channel whooping crane roosting habitat can be mechanically created and maintained but off-channel palustrine wetland roosting habitat is not a viable alternative to on-channel habitat due to a lack of potential restoration sites and costs associated with the creation of new wetland sites.
- Off-channel least tern and piping plover nesting habitat can be created and maintained using a variety of methods but on-channel nesting habitat is difficult to construct and erodes quickly.
- **Remaining Uncertainties** – relationship between channel width, vegetation, and flow and the combined effects of mechanical intervention and flow management actions; need for and mechanics of wet meadow and palustrine wetland management for whooping crane use.

Independent Scientific Advisory Committee (ISAC) Comments on the 2019 State of the Platte

The ISAC reviewed and discussed with GC members, the TAC, and the EDO the 2019 State of the Platte during the Program’s AMP Reporting Session in Omaha, NE in October 2019. During the AMP Reporting Session, the ISAC also discussed several “Deeper Dive Questions” (DDQs) developed by the EDO related to larger and more complex issues tied to development and implementation of a revised AMP for the Extension. Subsequently, the ISAC developed a set of recommendations in response to the draft State of the Platte and the DDQs which were included in the ISAC’s final 2019 report to the GC, presented to the GC in March 2020. The ISAC’s recommendations related to the Big Questions and the State of the Platte are summarized below in blue font. EDO responses to each recommendation are included below in green-shaded text boxes. For reference and further details, the full ISAC report to the GC is available online at www.PlatteRiverProgram.org.

From the ISAC: The following summarizes our recommendations on the 2019 State of the Platte (SOP) arranged by our report section and topic within each Section. Supporting text from our report is included for some recommendations to provide clarifying context.

Big Question Assessments for 2019 and First Increment.

BQ1: Clarify the use of two criteria: sandbar height 1.5’ above 1200 cfs (primary criterion); and sandbar height relative to peak flow.

Clarify the expected frequency of 15K cfs flows, their ecological benefits and flood risks.

EDO Response: Edits made in 2019 State of the Platte Report text.

BQ2: Figure 3 shows a large departure between observed versus predicted median Unobstructed Channel Width (UOCW) in 2016-2018. This departure is explained by, “there appears to be an additional driver (e.g., growing season

flows, etc.) for maintaining channel widths once channels are wide.” We recommend additional text briefly explaining how this departure may be addressed through the new 2-D modeling tool and other models presented during the AMP Reporting Session. Add text explaining how the new 2-D model can help to improve the ability to predict UOCW. The ISAC has suggestions for revising the structure of the decision tree model (see Section 2).

EDO Response: Edits made in 2019 State of the Platte Report text.

“Due to the degraded model performance from 2016-2018, the EDO has started to utilize machine learning random forest models that better incorporate hydrologic metrics, physical channel characteristics, and management activities to predict the cumulative effects of these metrics on channel attributes over time. Two-dimensional (2-D) modeling will parallel statistical modeling, providing improved predictions of physical channel characteristics such as inundated channel area, velocity, and shear stress over a range of flow conditions. Taken together, these modeling approaches will provide a more robust basis for development and testing of flow-habitat hypotheses during the Extension” (BQ2 – 2019 Assessment).

BQ3: We recommend BQ3 be carried forward to the Extension AMP for WC in parts of the system with sediment deficit, so as to maintain wide channels. Clarify that *you can* measure changes in bathymetry for a few miles downstream, but that it becomes more difficult as you move further downstream due to increasing uncertainty.

Recommendation: Ensure that statements of conclusions and management implications (pg. 15, SOP) are consistent with a one thumb up assessment or revise the assessment to reflect the reported high uncertainty of effectiveness of sediment augmentation to offset the deficit and halt channel degradation.

EDO Response: Edits made in 2019 State of the Platte Report text to make conclusions and management implications consistent with the one-thumb up assessment for this Big Question. This Big Question is currently under consideration for evaluation during the Extension as part of the revised Adaptive Management Plan (AMP).

BQ9: Whatever is decided for BQ9 in the 2019 assessment, we recommend that the authors be consistent in the evidence among BQs for assigning assessments.

To start the process of reconsidering BQ9, we reiterate our 2018 recommendation: *“The PRRIP should have clear expectations with respect to Program related benefits of proposed research on pallid sturgeon use of the Lower Platte River. This can be best accomplished in the short term by implementing the three tasks identified by Compass (2018, pg. 2) under The 2019 Decision: “What methods of reducing uncertainty should the Program pursue during the Extension to (a) better understand the role of the Platte in pallid recovery and (b) inform the connection between potential management alternatives and likely consequences on pallids?” For the longer term, the ISAC supports the 2030 decision step also described in Compass (2018, pg. 2): “What management actions should the Program undertake to best fulfill its obligations to pallid sturgeon in the Program’s Second Increment?”*

We also recommend that Lower Platte River pallid flow issues be embedded as a high-priority subset of the broader target flows topic when updating the AMP v2 during the First Increment Extension.

EDO Response: This Big Question and the associated issues are under consideration as part of the development of the revised AMP. As of March 2020, the Governance Committee (GC) is undergoing a process of discussing target flows and how, or if, to treat them as part of implementation of the revised AMP during the Extension. This includes discussion of the linkage between Program flow management actions and pallid sturgeon use and occurrence in the lower Platte River. The EDO recommends the GC consider these ISAC recommendations as they debate and decide on next steps for pallid sturgeon in the PRRIP.

BQ10: The Draft 2019 On-Channel Whooping Crane habitat assessment needs to better explain the evidence for changes in BQ rating to two-thumbs up, given that results shown in Figure 10 have changed little in the intervening 2 years and hypothesis S1c remains: “not yet answered - ongoing implementation, analysis and synthesis.”

EDO Response: Edits were made to the State of the Platte Report to indicate that an analysis of variance with post-hoc comparisons was used to compare distributions of annual maximum unobstructed channel width from 2007 – 2018 and indicated Program lands had significantly wider channel widths than non-Program lands since 2013. Pearson’s Rank Correlation showed a statistically significant increase in whooping crane use on Program Lands since 2016.

Additional Incremental Learning;

Least Terns and Piping Plovers. There is a need to more fully explain the scientific rationale for discontinuing efforts to construct in-river islands for tern and plover nesting and brood rearing habitat. Reference publications that provide such a rationale.

EDO Response: Edits made in 2019 State of the Platte Report text and citations added. Specific reasons were added, along with a primary reference source.

Appendix A. 2019 State of the Platte Priority Hypotheses Status Table

Because BQ8 P2 or BQ10 S1c are conclusively answered as two-thumbs up or down, why don't their priority hypotheses also receive a corresponding green-up or red-down triangle?

EDO Response:

P2 – this hypothesis has not been directly addressed through a study of plover forage (invertebrates) so as a distinct hypothesis it remains unanswered. However, there are no data suggesting declines in plover productivity or fitness related to forage. The PRRIP decided to forego further investigation into forage-related impacts for both terns and plovers.

S1c – this hypothesis has not been directly addressed via creation of additional wet meadow acres during the First Increment. Because Program monitoring does not suggest a selection preference for wet meadows by whooping cranes, this issue is not a high priority of concern for the Program during the Extension.

2007-2019 Big Question Assessments

To assist the GC with quickly evaluating the 2019 Big Question assessments, the icons in Table 1 are used to visually summarize the basic conclusion for each question. Thumbs up or down indicate a trend in the affirmative or negative and may point to the need to re-evaluate management actions based on collected data and analysis. The “unknown character” is used when there is not enough evidence to indicate a trend in either direction or more time is needed to collect appropriate data and conduct analyses. These icons are intended to provide the GC with a quick visual means to see where the Program stands each year in moving towards resolution of the Program’s most significant scientific questions as they relate to management decision-making.

Each Big Question assessment includes an indicator of the “test results” for relevant priority hypothesis. Conclusions about the hypothesis are indicated as one of the following categories:

-  Hypothesis answered conclusively – affirmed.
-  Hypothesis answered conclusively – rejected.
-  Hypothesis not yet answered – ongoing implementation, analysis, and synthesis.
-  Not currently being addressed through implementation of the AMP and related data analysis and synthesis.

See **Appendix A** for a more detailed status report for each priority hypothesis in the AMP.

TABLE 1. Quick Reference legend explaining icons used to assess Big Questions.

Icon	Trend or Answer Explained by Icon
	<ul style="list-style-type: none"> • Big Question and underlying hypotheses answered conclusively in the affirmative • Foundational documents, analysis, and other references on which this assessment is based have undergone peer review through the PRRIP peer review process and/or publication in refereed journals • Governance Committee should consider adjustments to decisions related to PRRIP management actions
	<ul style="list-style-type: none"> • Affirmative answer or trend, but Big Question and underlying hypotheses NOT answered conclusively • Assessment can be based on draft documents and analysis, but peer review and/or publication may be pending • To the extent possible, consider what information is necessary to change this designation
	<ul style="list-style-type: none"> • Evidence thus far is inconclusive; no affirmative or negative answer/trend to Big Question and underlying hypotheses • Assessment can be based on draft documents and analysis, but peer review and/or publication may be pending • To the extent possible, consider what information is necessary to change this designation
	<ul style="list-style-type: none"> • Negative answer or trend, but Big Question and underlying hypotheses NOT answered conclusively • Assessment can be based on draft documents and analysis, but peer review and/or publication may be pending • To the extent possible, consider what information is necessary to change this designation
	<ul style="list-style-type: none"> • Big Question and underlying hypotheses answered conclusively in the negative • Foundational documents, analysis, and other references on which this assessment is based have undergone peer review through the PRRIP peer review process and/or publication in refereed journals • Governance Committee should consider adjustments to decisions related to PRRIP management actions

TABLE 2. Big Question assessments, PRRIP First Increment (2007-2019).

PRRIP Big Question	First Increment Assessment	Basis for Assessment
<u>Implementation</u> – Program Management Actions and Habitat		
1. Will implementation of SDHF produce suitable tern and plover riverine nesting habitat on an annual or near-annual basis?		<u>Conclusively answered.</u>
2. Will implementation of SDHF produce and/or maintain suitable whooping crane riverine roosting habitat on an annual or near-annual basis?		<u>Conclusively answered.</u>
3. Is sediment augmentation necessary for the creation and/or maintenance of suitable riverine tern, plover, and whooping crane habitat?		Trending positive and certainty about the sediment deficit in the south channel above the Overton bridge; uncertainty about the role of that deficit in habitat creation and maintenance in the rest of the AHR. This Big Question will either be retained in its current form or revised and addressed through implementation of AMP Version 2.0 during the Extension.
4. Are mechanical channel alterations (channel widening and flow consolidation) necessary for the creation and/or maintenance of suitable riverine tern, plover, and whooping crane habitat?		<u>Conclusively answered.</u>
<u>Effectiveness</u> – Habitat and Target Species Response		
5. Do whooping cranes select suitable riverine roosting habitat in proportions equal to its availability?		<u>Conclusively answered.</u>
6. Does availability of suitable nesting habitat limit tern and plover use and reproductive success on the central Platte River?		<u>Conclusively answered.</u>
7. Are both suitable in-channel and off-channel nesting habitats required to maintain central Platte River tern and plover populations?		<u>Conclusively answered.</u>
8. Does forage availability limit tern and plover productivity on the central Platte River?		<u>Conclusively answered.</u>
9. Do Program flow management actions in the central Platte River avoid adverse impacts to pallid sturgeon in the lower Platte River?		This Big Question will either be retained in its current form or revised and addressed through implementation of AMP Version 2.0 during the Extension.
10. Do Program management actions in the central Platte River cumulatively 1) produce detectable changes in the physical environment (i.e. habitat) and 2) result in a detectable increase in tern, plover, and whooping crane use of the Associated Habitats?	<p style="text-align: center;"><u>LTPP Off-Channel</u></p> <p>Habitat  Species Response </p> <p style="text-align: center;"><u>WC On-Channel</u></p> <p>Habitat  Species Response </p>	<u>Conclusively answered.</u>



Big Question #1

Will implementation of Short-Duration High Flow releases produce suitable tern and plover riverine nesting habitat on an annual or near-annual basis?




Figure 1. First Increment peak flow event magnitudes and volumes in relation to SDHF. Seven events (2008, 2010, 2011, 2013, 2015, and 2016) exceeded SDHF magnitude and duration and only the extreme event in 2015 produced any suitably high sandbar nesting habitat (see Figure 2).

2019 Assessment



- Observational studies of natural high flow events since 2007 have provided sufficient data to test the hypothesis that SDHF releases will create suitably-high sandbars.
- The minimum sandbar height suitability criterion for nest initiation is ≥ 1.5 ft above 1,200 cfs river stage. Full SDHF magnitude of 8,000 cfs is not sufficient to create sandbars exceeding this height suitability criterion.
- Sandbars created by SDHF releases will be inundated during the nesting season in most years.
- Peak flow magnitudes of 15,000 cfs will produce sandbars meeting the minimum height criterion. However, this flow magnitude occurs infrequently (10-year return interval), the quantity of suitably high sandbar area would be well below the AMP objective of 10 acres per river mile, and sandbars at this height would be inundated during the nesting season in many years.




Figure 2. Distribution of emergent sandbar area produced during the 2015 peak flow event in the portion of the AHR downstream of Kearney. The 15,000 cfs event produced 43 cumulative acres (line) of sandbar habitat exceeding the minimum height suitability criterion of 1.5 ft above 1,200 cfs stage. Median height of bars was 1.6 ft above 1,200 cfs stage.



What the science says in 2019:

- The original analysis of SDHF performance assumed sandbars build to the water surface during peak flow events. The median height of sandbars formed during natural high flow events in 2010, 2011, 2014, and 2015 was 1.2 – 2.3 ft below peak stage. At this relative height, sandbars created in 2010, 2011 and 2014 were well below the Program’s minimum height criterion for nest initiation of ≥1.5 ft above 1,200 cfs stage.¹
- A natural high flow event of 15,000 cfs in 2015 produced sandbars exceeding the minimum height criterion. The median height of sandbars formed in 2015 was 1.6 ft above 1,200 cfs stage (Figure 2) with approximately 43 acres of mid-channel bar area ≥1.5 ft above 1,200 cfs stage were present in the portion of the AHR downstream of Kearney in November of 2015 (Figure 2). This equates to 0.8 acres per river mile.

We estimate with confidence that:

- SDHF magnitude of 5,000 to 8,000 cfs for a duration of three days at peak would not be sufficiently long to mobilize the bed and produce suitable sandbar nesting habitat.
- Sandbars created by a full SDHF magnitude of 8,000 cfs would be 0.5 – 1.0 ft lower than the minimum height criterion and would be inundated at flows experienced in the AHR during most nesting seasons.
- Peak flow magnitudes of 15,000 cfs occur infrequently (10-year return interval) but will produce sandbars exceeding the minimum height criterion given sufficiently long duration at peak.
- Even at a discharge magnitude of 15,000 cfs, total suitable sandbar area would be well below the AMP objective of 10 acres per river mile. Additionally, sandbars at the 2015 median height would be inundated during the nesting season in many years limiting the potential for successful reproduction.


Answering BQ #1 during the First Increment

- Six tern/plover habitat synthesis chapters and associated publications serve as the best source for synthesized reference data for this question. Those chapters have been peer reviewed and accepted by the Governance Committee² and three of those chapters have been published.
- Geomorphic and species monitoring data collected through 2018 are consistent with and support the analyses and conclusions presented in the synthesis chapters and associated publications.

Management Implications:

- Big Question #1 has been answered with a definitive “two thumbs down.” The Governance Committee completed the final “Adjust” stage of adaptive management and decided to maintain 10 acres of on-channel moving complex

approach (MCA) islands and to create an additional 60 acres of off-channel nesting habitat.



Priority Hypothesis	Results
Flow #1	▼
<p>Relationship to Big Question #1 Based upon the SedVeg model and associated assumptions in the FSM management strategy, it is hypothesized that under a balanced sediment budget, flows of 5,000 to 8,000 cfs magnitude for three days (SDHF) will build sandbars to an elevation that is suitable for tern and plover nesting. The Program’s minimum height suitability criterion is 1.5 ft above 1,200 cfs river stage and represents the minimum height thought necessary for nest initiation.</p>	



Big Question #2

Will implementation of Short-Duration High Flow releases produce and/or maintain suitable whooping crane riverine roosting habitat on an annual or near-annual basis?

Channels with unobstructed channel widths ≥ 650 ft and unforested corridor widths $\geq 1,100$ ft are highly suitable for whooping crane roosting. See Big Question 5.




Figure 3. Observed versus predicted median unobstructed channels widths (UOCW) in the AHR during the period of 2007-2018. The UOCW model provides good predictive capacity for evaluating the efficacy of SDHF releases prior to 2016⁵. Poor predictive performance in 2016-2018 indicates there is an additional driver that is maintaining UOCW during that period despite low 40-day mean peak discharge magnitudes.

2019 Assessment



- Mature phragmites plants or plant patches have a very low probability of being eroded at the highest flow magnitudes and velocities observed in the AHR. An herbicide control program is ongoing.³
- Program analyses strongly support the assertion of a positive relationship between peak flow magnitude and unobstructed channel width (UOCW) in the AHR during the period of 2007 - 2015. During this period, 40-day mean peak discharge was the best hydrologic predictor of UOCW.
- The comparatively short duration and low volume of SDHF limits the predicted increase in UOCW to ≤ 12 ft. SDHF magnitude and duration is not sufficient to create and maintain UOCWs that are suitable for whooping crane roosting.
- 40-day mean peak discharge is not a good predictor of UOCW in 2016-2018 when UOCW remained high despite low 40-day mean peaks.
- We hypothesize that UOCW remained high in 2016-2018 due to channel inundation during the growing season which prevented vegetation germination.
- The short duration SDHF is insufficient to inundate the channel long enough to prevent vegetation germination. SDHF would not maintain UOCWs that are suitable for whooping crane roosting.
- Due to the degraded model performance from 2016-2018, the EDO has started to utilize machine learning random forest models that better incorporate hydrologic metrics, physical channel characteristics, and management activities to predict the cumulative effects of these metrics on channel attributes over time. Two-dimensional (2-D) modeling will parallel statistical modeling, providing improved predictions of physical channel characteristics such as inundated channel area, velocity, and shear stress over a range of flow conditions. Taken together, these modeling approaches will provide a more robust basis for development and testing of flow-habitat hypotheses during the Extension.



What the science says in 2019:

- Phragmites occurrence and percent cover declined significantly during the period of 2009-2012 and were slightly increasing to stable in more recent years. The reduction is positively correlated with herbicide application and not correlated with peak flow magnitude or inundation duration.⁴
- 40-day mean peak discharge was the best hydrologic predictor of UOCW in the AHR during the period of 2007-2015. Other metrics useful in predicting UOCW included bankfull wetted width, median bed material grain size, and whether spraying or disking occurred.⁵
- During the period of 2016-2018 predictive performance is poor as UOCW remained high despite low 40-day mean peak discharges. We hypothesize that channel inundation during the growing season maintained UOCW by preventing vegetation establishment.

We estimate with confidence that:


- Implementation of a three- to five-day SDHF will have a minimal influence on UOCW in the AHR during years SDHF releases could have been made would have resulted in ≤ 12 ft increase in the maximum unobstructed width within a year, but the effect would not carry over to subsequent years due to vegetation encroachment.
- The limited benefit of SDHF is not sufficient to produce suitably wide UOCWs during dry years.
- During wet years, flow releases are not necessary to produce suitably wide UOCWs.
- Implementation of disking and herbicide increases UOCW by an average of 126 ft, producing suitably wide UOCW.
- Assuming inundation maintained UOCW after 2016, SDHF duration is insufficient to inundate the channel long enough to prevent vegetation germination.

Answering BQ #2 during the First Increment

- The Program has published directed scour research which serves as the best source for synthesized reference data for phragmites scour resistance.³
- The Program’s whooping crane data synthesis chapters and associated publications serve as the best source for synthesized reference data for the relationship between SDHF and unvegetated channel width. Those chapters have been peer reviewed and accepted by the Governance Committee⁵ and have subsequently been published¹¹.
- The Program is developing new modeling tools that incorporate the effects of both peak flows and channel inundation on UOCW. These tools will be used by the Program during the Extension to explore alternative flow management actions including low-magnitude long-duration releases to prevent germination and maintain UOCW.

Management Implications:

- Implementation of SDHF releases as currently envisioned is highly unlikely to create and/or maintain suitably wide UOCWs for whooping cranes.
- Implementation of disking and herbicide application at Program habitat complexes will create and maintain suitably wide UOCWs for whooping cranes.
- Low-magnitude longer-duration flow releases may help maintain UOCW. This will be explored during the Extension.





Big Question #3

Is sediment augmentation necessary for the creation and/or maintenance of suitable riverine tern, plover and whooping crane habitat?

2019 Assessment



- The south channel reach from the J2 Return to the Overton bridge is incising and narrowing due to the sediment imbalance from clear water hydropower returns. Downstream from Overton, the large degree of spatial and temporal variability in channel form makes it difficult to draw conclusions about sediment balance.
- South channel degradation began following the construction of the J-2 Return in the early 1940s has resulted in a portion of that reach transitioning from a wide braided planform to a narrow wandering planform, which is less suitable for use by the Program’s target species.

- Augmentation of sediment in the south channel is necessary to slow incision and narrowing and prevent degradation from progressing downstream past the Overton bridge. While necessary for maintaining suitable channel widths for whooping cranes, it is not necessary for increasing productivity of terns and plovers within the AHR.
- It will be challenging to measure the effectiveness of augmentation given the desired beneficial effect is slowing and ultimately halting a long-term degradational trend to prevent degradation downstream of the Overton bridge. As such, we believe it will take another 5-7 years of sediment augmentation, analyses, and observations to determine the effectiveness of sediment augmentation in halting the downstream movement of incision of the channel bed.
- This big question will continue to be assessed during the First Increment Extension.




Figure 3. 2009 to 2014 longitudinal profile and volume change for the reach from J2 Return to Overton bridge. Volume change in the reach immediately downstream of the J-2 Return is caused by channel incision. Further downstream, volume change is due primarily to channel widening.



What the science says in 2019:

- Sediment transport modeling indicates a mean annual sediment deficit of 55,000 tons in the south channel segment extending from the J2 Return downstream to the Overton bridge, ranging from 0 tons in dry years to >100,000 tons in wet years.⁶
- Between 2009 and 2014, that reach lost an average of 159,000 tons of sediment annually due to incision and lateral erosion of banks.⁷
- Incision and associated reduction in channel slope were greatest immediately downstream of the J2 Return and was negligible at the Overton bridge (Figure 3).
- Full-scale sediment augmentation will not be 100% efficient. A proportion (~10%) of the augmentation material will either be too coarse to be mobilized from the augmentation site or so fine that it is rapidly transported out of the reach.

We estimate with confidence that:

- Observed incision and narrowing and associated planform change in the south channel result in a channel configuration that is not suitable for use by the Program’s target species.
- In absence of augmentation to offset the south channel deficit, incision and narrowing would progress downstream past the Overton bridge and negatively affect habitat suitability at the Program’s Cottonwood Ranch complex.
- Early indications are that augmentation of 60,000 to 80,000 tons of sand annually downstream of the J2 return appears to be sufficient to allow the Program to evaluate augmentation efficiency.
- Measuring augmentation effectiveness will require assessment of changes in channel slope, volume, width, and bed material. It may be challenging to quantify beneficial effects for whooping cranes, but changes in channel form and configuration should be more evident.


Answering BQ #3 during the First Increment

- The existence and negative impacts of a sediment deficit downstream of the J2 Return has been well documented by the Program and others.
- The effectiveness of sediment augmentation in offsetting the deficit and halting degradation is not known, but preliminary data indicates it will be effective.
- Full scale operations began in the fall of 2017 and it is anticipated that five to seven years of implementation and response monitoring will be necessary to assess augmentation efficiency and effectiveness.

Management Implications:

- If the south channel sediment deficit persists, incision and narrowing will progress downstream past the Overton bridge, negatively influencing whooping crane habitat suitability in an increasingly larger portion of the AHR.

- Early indications are that full scale sediment augmentation will be effective in halting the long-term trend of incision and narrowing. The beneficial effects of augmentation need to be assessed through five to seven years of implementation and effectiveness monitoring during the First Increment Extension that will include biannual bathymetric LiDAR collection and analysis.
- Between baseline conditions and the end of the second year of augmentation, a detectable channel response is observed in the upper 5 miles of the augmentation area with respect to slope and channel width. On average, minimum bed elevations have increased 0.9 ft in this reach; 1.2 ft in the upper 2 miles of the reach. Channel widths have increased an average of 70 feet during this period between the J2 return and Overton Bridge. An average width increase of 90 feet was observed in the upper 5 miles and the upper 2 miles have increased by 160 ft.



Priority Hypothesis	Results
Sediment #1	●
Relationship to Big Question #3	
Based on the SedVeg model and associated assumptions in the FSM management strategy, it is hypothesized that eliminating the existing sediment deficit through sediment augmentation is necessary to reduce channel narrowing and incision, contribute to channel widening, and increase the sustainability of a braided channel morphology.	



Big Question #4

Are mechanical channel alterations (channel widening and flow consolidation) necessary for the creation and/or maintenance of suitable riverine tern, plover, and whooping crane habitat?

2019 Assessment



- Peak flows in the AHR are generally not sufficient to remove mature woody vegetation or erosion-resistant species like phragmites.
- Mechanical clearing and leveling are necessary to create suitable channel configurations and facilitate channel adjustments to changes in flow and sediment.
- Ongoing mechanical management actions like herbicide application and disking are necessary to maintain suitably wide unobstructed channel widths (UOCWs) for target species.
- Flow consolidation, a mechanical management action which consists of mechanically confining 90% of total river flow into a single channel, may support the maintenance of suitable UOCWs but is not implementable due to regulatory and legal impediments.




Figure 4. Predicted probability of a transect measuring ≥ 650 ft in unobstructed channel width (suitable for whooping cranes) in relation to 40-day peak discharge at transects with (blue solid line) or without (red dashed line) mechanical management actions in the AHR from 2007 to 2018. Shaded regions indicate 95% confidence intervals. Disking and herbicide application provide a significantly greater probability of having channels with more than 650 ft unobstructed channel widths.



What the science says in 2019:

- Phragmites is extremely erosion resistant and SDHF flow depths and velocities are only sufficient to scour the very weakest individual plants. Ability to scour woody vegetation also decreases dramatically in the year following seed germination.⁸
- Locations that are mechanically maintained through herbicide application and disking have a significantly higher probability of being suitably wide for whooping crane roosting (Figure 4).

We estimate with confidence that:

- Mechanical clearing, leveling, and channel widening are necessary to create suitably wide channels at Program habitat complexes.
- Herbicide application and disking are necessary at Program habitat complexes in most years to maintain suitably wide UOCWs.
- The beneficial effects of mechanical management actions are largely limited to the locations where they are implemented. They do not provide system-scale beneficial effects.

Answering BQ #4 during the First Increment

- The Program has published directed scour research which serves as the best source for synthesized reference data for phragmites scour resistance.³
- The Program’s whooping crane data synthesis chapters are the best source for synthesized reference data for the relationship between mechanical actions and unvegetated channel width. Those chapters have been peer reviewed and accepted by the Governance Committee.⁵

Management Implications:

- It was originally hypothesized that mechanical actions were necessary to create desired channel configurations that would subsequently be maintained through Short Duration High Flow releases. SDHF has been shown to be ineffective at creating suitable tern and plover nesting habitat and maintaining suitable channel widths for whooping cranes. Accordingly, ongoing mechanical maintenance will be necessary to provide nesting habitat and maintain suitable UOCWs at Program habitat complexes.
- Due to regulatory and legal issues flow consolidation has been abandoned as a potential Program management action.

Priority Hypothesis

Mechanical #2

Results



Relationship to Big Question #4

Based on the SedVeg model and associated assumptions in the FSM management strategy, it is hypothesized that designed mechanical channel alterations like flow consolidation, mechanical clearing and leveling of islands, channel widening, and vegetation clearing from banks are needed to accelerate the creation of, and/or to maintain suitably wide braided channels in the AHR.





Big Question #5

Do whooping cranes select riverine roosting habitat in proportions equal to its availability?




Figure 5. Distribution of unobstructed channel width (UOCW) at use (n=235) and available riverine roost locations in the Associated Habitat Reach (AHR). Use locations were selected disproportionately more than availability above 689 feet of UOCW¹¹. As such the Program has agreed to manage for 650 ft channels for whooping crane roosting habitat on the central Platte River. Density curves are represented as dashed (use) or dotted (available) lines.

2019 Assessment



- Results of habitat selection analyses within the AHR and throughout the Great Plains indicate whooping cranes select unobstructed channel widths of ~650 feet and unforested corridor widths of ~1,100 disproportionately to availability.^{9,10,11}




Figure 6. Distribution of nearest forest (NF) at use (n=235) and available riverine roost locations in the Associated Habitat Reach (AHR). Use locations were selected disproportionately more than availability above 594 feet of NF¹¹. As such the Program has agreed to manage for total unforested corridor widths of ~1,100 ft (suitable NF roughly multiplied by 2) which would be highly favorable for whooping cranes roosting on the central Platte River. Density curves are represented as dashed (use) or dotted (available) lines.



What the science says in 2019:

- First Increment habitat management efforts implemented by the Program to date include, but are not limited to, tree removal and bank line diking to increase unobstructed view widths, channel diking and widening to increase unobstructed channel widths, and flow releases and sediment augmentation to test hypotheses related to increasing river braiding and areas of suitable depth for whooping crane roosting.

We estimate with confidence that:

- Whooping cranes select unobstructed channel widths of ≥ 689 feet and unforested corridor widths of $\geq 1,188$ feet disproportionately to availability.¹¹

Answering BQ #5 during the First Increment

- Detailed habitat selection analyses have been completed and have undergone the Program's independent third-party peer review^{9,10} and publication¹¹. The Program accepted the whooping crane habitat synthesis chapters and the WEST whooping crane report and peer reviews as final.

Management Implications:

- Based on findings of habitat selection analyses, the Program has agreed to continue to manage to provide unobstructed channel widths that are ≥ 650 ft and unforested corridor widths that are $\geq 1,100$ ft.



Priority Hypothesis	Results
WC 3	▼
Relationship to Big Question #5	
It is hypothesized that whooping crane use is related to habitat suitability values as defined in Land Plan Table 1.	

WC 3

Results



Relationship to Big Question #5

It is hypothesized that whooping crane use is related to habitat suitability values as defined in Land Plan Table 1.

Photo Credit: Abby Jensen



Big Question #6

Does availability of suitable nesting habitat limit tern and plover use and reproductive success on the central Platte River?

2019 Assessment

- Long-term monitoring and data analyses indicate there is a strong positive correlation between Program-defined suitable off-channel **nesting** habitat and tern and plover breeding pair counts within the AHR.^{12,13} During the Program's First Increment, the tern and plover populations on the central Platte River have increased significantly (Pearson's Rank Correlation test $\alpha \leq 0.05$) and proportionately to increases in habitat availability.




Figure 7. Tern (dotted line; top) and plover (dashed line; top) total breeding pair counts and Program (94 ac) and non-Program (84 ac) habitat availability (solid blue line) based on Program habitat availability assessments and tern (bottom left) and plover (bottom right) reproductive success as compared to the Lutey (2002) objectives, 2007-2019. Increased off-channel habitat availability and high reproductive success within the AHR are believed to be responsible for the increases tern and plover populations on the central Platte River.



What the science says in 2019:

- The Program and its partners have created in-channel (sandbars) and off-channel (sandpits) nesting habitat to evaluate hypothesized relationships between habitat availability and tern and plover use and productivity within the Program Associated Habitat Area. The Program has created and maintained ~90 acres of off-channel and ~65 acres of in-channel nesting habitat for terns and plovers.¹¹ In addition, Program partners and mining operations have constructed and/or managed ~60 acres of off-channel and ~25 acres of in-channel nesting habitat.
- Numbers of tern and plover breeding pairs have increased 4-fold within the AHR since 2001 while increases of similar magnitude have not been observed on Lake McConaughy (Dave Zorn, personal communication) or the Missouri River (<http://moriverrecovery.usace.army.mil/mrrp/f?p=136:6:0::NO>); although recent increases have been observed on the Missouri River. While overall numbers of tern and plover breeding pairs within the AHR have increased significantly (Pearson's Rank Correlation test $\alpha \leq 0.05$), habitat availability and use of non-Program habitat has remained steady¹¹. We have observed a high, positive correlation between tern and plover breeding pair counts and habitat availability. Program data also indicate breeding pair counts increase at a similar rate as habitat availability.
- Piping plover reproductive success, as measured by fledglings/breeding pair, has recently declined below the Lutey (2002) objective for maintaining stable to increasing populations within the AHR. Least tern fledge ratios have declined but remain above the Lutey (2002) objective.

We estimate with confidence that:

- There is a high correlation between habitat availability and breeding pair counts and as the Program increases suitable off-channel nesting habitat, numbers of tern and plover breeding pairs within the AHR will increase until habitat availability exceeds population demands.

Answering BQ #6 during the First Increment

- Tern and plover data collected to date and published in the 2015 Breeding Pair publication¹² and 2017 tern and plover habitat selection publication¹⁴ serves as the best source data for this question.
- The 2018 Tern and Plover Monitoring and Research Report¹¹ has also been reviewed and accepted by the Program and serves as additional evidence of the ongoing increasing trend in tern and plover use of the AHR.

Management Implications:

- Based on results of Program analyses, the Program has agreed to increase off-channel habitat availability by 60 acres.



Priority Hypothesis	Results
T1	▲
P1	▲
Relationship to Big Question #6 It is hypothesized that when in-channel (sandbars) and off-channel (sandpits) nesting habitat availability increase, tern and plover use and productivity will increase (i.e., habitat is limiting).	





Big Question #7

Are both suitable in-channel and off-channel nesting habitats required to maintain central Platte River tern and plover populations?

2019 Assessment



- Long-term monitoring and data analyses indicate a combination of both in-channel and off-channel *nesting* habitats are not necessary to maintain the central Platte River population of terns and plovers. During the Program's First Increment the increase in tern and plover populations on the central Platte River is the result of use and productivity at off-channel nesting habitats.¹¹ River survey and observational data, however, indicate the river is a valuable source of *forage* for both species as forage availability appears to be lower on off-channel habitats.¹⁵




Figure 7. Annual tern (left plot) and plover (right plot) total (solid line), riverine (dotted line), and sandpit breeding pair counts (dashed line), 2001-2019. Trends in use of the AHR represent significant increases in tern and plover breeding pair counts during 2001-2019 with the most substantial increases occurring since inception of the Program. Reproductive success on and use of sandpits is responsible for the increase.



What the science says in 2019:

- The Program and its partners created in-channel (sandbars) and off-channel (sandpits) nesting habitat to evaluate hypothesized relationships between in- and off-channel habitat availability and selection of terns and plovers. Early Program efforts largely focused on off-channel nesting sites as flows and permitting challenges precluded construction of in-channel nesting islands. Program efforts in recent years were directed at maintaining off-channel nesting habitat and constructing and maintaining suitable in-channel habitat.
- The creation and maintenance of off-channel nesting habitat resulted in substantial use and productivity since 2001. During this same timeframe, in-channel habitat availability and tern and plover nesting and productivity have been sporadic and thus has not contributed to the maintenance of the central Platte River populations. Despite the limited use and productivity of in-channel nesting habitat, we observed significant increases in the numbers of tern and plover breeding pairs within the AHR from 2001-2019.¹¹
- Since 2001, breeding pair counts for terns increased over 4-fold (21 to 95) while plover counts also increased over 4-fold (10 to 45); both of which represent significant increases.¹¹ Though populations of both species increased during this timeframe, increases of similar magnitude have not been observed on Lake McConaughy (Dave Zorn, personal communication) or the Missouri River (<https://usace.contentdm.oclc.org/utills/getfile/collection/p16021coll3/id/825>); though recent increases have been observed on the Missouri River.
- Efforts to create and maintain suitable in-channel nesting habitat have necessarily been extensive. Though in-channel nesting habitat has contributed little to the reproductive success of both populations, ephemeral islands and river channels appear to provide an important source of forage for both terns and plovers.

We estimate with confidence that:

- Managed off-channel nesting habitat is necessary to maintain the presence of central Platte River tern and plover populations.
- Although an important forage source, direct maintenance of in-channel nesting habitat is not necessary to maintain tern and plover populations within the AHR.

Answering BQ #7 during the First Increment

- Tern and plover monitoring data collected to date and the 2015 Breeding Pair publication¹² serve as the best source data for this question and indicate use of off-channel habitat resulted in increases in breeding pair counts and productivity within the AHR.
- The 2018 Tern and Plover Monitoring and Research Report¹¹ has also been reviewed and accepted by the Program and serves as additional evidence of the ongoing increasing trend in tern and plover use of the AHR attributable to use of and productivity on off-channel sites.

Management Implications:

- Based on results of Program analyses, the Program has agreed to increase off-channel habitat availability by 60 acres while creating and maintaining 10 acres of MCA islands in the channel.



Priority Hypothesis	Results
TP1	▼
<p>Relationship to Big Question #7 It is hypothesized that ephemeral, in-channel nesting islands (sandbars) are needed for long-term nesting success of terns and plovers on the central Platte and when available, terns and plovers will select sandbars over sandpits for nesting. It is also hypothesized that tern and plover nesting is more successful on in-channel than off-channel habitat which could eliminate the need to maintain off-channel habitat.</p>	



Big Question #8

Does forage availability limit tern and plover productivity on the central Platte River?




Figure 9. Cumulative proportion of least tern broods that fledged at various flows during the brood rearing season, 2001-2019 (top) and annual tern fledge ratio versus median summer flows, 2001-2019 (bottom). Blue vertical lines represent the hypothesized 800 cfs target flow and the horizontal red dashed line represents the Lutey (2002) fledge ratio objective for maintaining a stable to growing population on the central Platte River.

2019 Assessment



- Analyses of flow versus productivity¹⁶ indicate there is no relationship between flow and tern productivity and we suspect analyses of data linking forage availability or flow and plover productivity would yield similar results. Given tern and plover productivity is high and most confirmed mortalities have been attributed to adverse weather and predation, there is no evidence the forage base along the central Platte River limits tern and plover productivity. Further evaluations would involve capturing and weighing tern and plover chicks on multiple occasions to establish a more direct link between growth rates and forage abundance; however, Program stakeholders decided these additional expenses, efforts, and risk of injury to chicks are not warranted.





What the science says in 2019:

- Detailed analyses have been completed and the resulting manuscript has been published. In the publication, we synthesize independent sets of data and found no relationship between tern productivity and flow during the nesting and brood rearing season.^{15,16}
- Given the high levels of productivity observed on the central Platte River, it is unlikely flow, and thus forage fish abundance, limits tern productivity. We were unable to establish the hypothesized link between flow and productivity and have used results of our retrospective analyses to definitively answer this Big Question.
- Further evaluations of BQ #8 would likely entail system-wide, intensive, summer-long forage sampling, tern and plover behavioral studies, and potentially capturing and weighing chicks on multiple occasions to attempt to establish relationships between forage abundance, flow, productivity, and long-term survival. Program stakeholders previously indicated additional expenses, efforts, and risk of injury to chicks are not warranted as it appears forage abundance and reproductive success are adequately high to support central Platte River tern and plover populations.

We estimate with confidence that:

- Forage availability does not limit tern and plover productivity on the central Platte River.

Answering BQ #8 during the First Increment

- The forage fish manuscript¹⁵ serves as the best source for synthesized reference data for this question. The results of these analyses indicate flow, and by inference forage availability, does not limit tern and plover productivity within the AHR. Program staff considered results of these analyses to be sufficient evidence to change the assessment for this Big Question to two thumbs down in 2016.
- A similar synthesis of data could be developed for plovers; however, given results of the Foraging Habits Study and high levels of productivity observed to date, there is a complete lack of evidence forage abundance limits plover productivity.

Management Implications:

- Data analysis and synthesis do not support Program summer flow releases to maintain the 800 cfs target.

Priority Hypothesis	Results
T2	Red Triangle
P2	Yellow Circle

Relationship to Big Question #8
It is hypothesized that availability of fish for terns and invertebrates for plovers limits productivity of both species, especially when flows are below 800 cfs during the nesting season (May through August).



Big Question #9

Do Program flow management actions in the central Platte River avoid adverse impacts to pallid sturgeon in the lower Platte River?



2019 Assessment



- The Program's Lower Platte River Stage Change Study¹⁶ analyses concluded central Platte River flow management actions are likely to avoid adverse impacts to pallid sturgeon in the lower Platte River because the relative change in habitat due to Program water management activities would be small to undetectable.
- Based on results of the Stage Change Study, any potential impacts could be avoided through development of operational rules that prohibit Program diversions when lower Platte River discharges fall below 4,000 cfs.
- Thus, EDO followed the established process to assess this Big Question as conclusively answered with two thumbs up in 2014. In 2016, the Service concluded they "do not support two thumbs up at this time for Big Question 9 because of lingering uncertainties."
- There remains substantial scientific uncertainty about pallid sturgeon use of the lower Platte, though there has been substantial new learning about pallid sturgeon and their use of lower Platte (evidence they are there all year, spawning ground, larval drift, etc.) since the Stage Change Study was completed. At the time of Stage Change Study, the primary issue was use of the lower Platte River by a small number of adult fish.
- A 2018 EDO white paper identified substantial uncertainties with pallid sturgeon use of the lower Platte River and also uncertainties regarding the ability of the Program to conduct research alone or in partnership with entities such as the Nebraska Game & Parks Commission, the University of Nebraska-Lincoln, and/or the U.S. Geological Survey that would render useful data for GC decision-making.
- In March 2019, the GC directed the EDO to integrate pallid sturgeon uncertainties into the process for updating/revising the AMP for the Extension. That process began in May 2019.

What the science says in 2019:

- The general conclusion of the Program’s Final Stage Change Study¹⁶ is that Program water management activities will not result in measurable changes on flows in the lower Platte River and thus will result in little change to the amount of habitat available to pallid sturgeon.
- However, given that short-term connectivity could be problematic under certain, but infrequent, hydrological conditions, and assuming the biological significance of habitat connectivity for pallid sturgeon above 4,000 cfs, results of the stage change study could be used by the Program to implement proactive measures (e.g. altering excess-to-target-flow diversion timing or duration) to prevent potential negative impacts on habitat connectivity.

We estimate with confidence that:


- Flow diversions or releases by the Program would result in very small and undetectable changes in stage in the lower Platte River.
- As identified in the stage change study, these stage changes reside in the noise of gage error on the lower Platte River and thus will not result in a measurable change in lower Platte River stage.
- By extension, flow management actions that will not result in a measurable change in stage in the lower Platte River will not result in significant adverse effects on pallid sturgeon.

Answering BQ #9 during the First Increment:

- This question is not likely to be answered until the First Increment Extension. The GC began a facilitated Pallid Sturgeon Process in 2017 to help guide activities that will keep the question open until sometime during the potential First Increment Extension. That effort was integrated into the development of AMP Version 2.0 for the Program beginning in May 2019.

Management Implications:

- The primary Program water management actions that are hypothesized to result in flow and fish impacts in the lower Platte River are short-duration high flows (SDHF), target flow releases, and diverting target flow excesses.
- The Program is undergoing a process to develop flow management actions for the potential First Increment Extension.
- Central Platte River flow releases or diversions that could plausibly be detected in the lower Platte River during the potential First Increment Extension are not anticipated.



Priority Hypothesis	Results
PS2	
Relationship to Big Question #9 It is hypothesized that Program water management actions, such as diverting excesses to target flows for retimed release, will result in a measurable change in stage in the lower Platte River and thus affect pallid sturgeon habitat suitability.	

Credit: Sam Stukel

Big Question #10

Do Program management actions in the central Platte River cumulatively produce detectable changes in the physical environment (i.e. habitat) that are associated with a detectable increase in tern, plover and whooping crane use of the Associated Habitats?

The Program implements both on- and off-channel habitat creation and maintenance for the target species. The BQ 10 assessment will focus on the habitat creation strategy that has been most effective for each species. In the case of least terns and piping plovers, off-channel habitat has been most effective. In the case of whooping cranes, on-channel habitat has been most effective.

2019 Assessment for Least Tern and Piping Plover Off-Channel (OCSW) Habitat

- There is a strong positive correlation between Program-defined suitable nesting habitat and tern and plover breeding pair counts within the AHR. See BQ 6 Assessment.¹¹
- During the First Increment, tern and plover populations on the central Platte River have increased significantly (Pearson's Rank Correlation test $\alpha \leq 0.05$) and proportionately to increases in habitat availability due to Program off-channel habitat creation efforts. See BQ 6 Assessment.¹¹

Habitat: 🍑🍑

Species Response: 🍑🍑

2019 Assessment for On-Channel Whooping Crane Habitat

- Maximum unobstructed channel widths (important whooping crane habitat metric) have been significantly wider (analysis of variance with yearly post-hoc comparisons adjusted- $\alpha \leq 0.05$) on Program lands since 2013. See Figure 10.

Habitat: 🍑🍑

Species Response: 🍑🍑




Figure 10. Unobstructed channel width (UOCW), an important whooping crane habitat suitability metric, have increased with Program management actions (bars) and has been wider on Program-managed lands (solid line) since 2011 and substantially wider since 2013. Program lands are generally now more suitable for whooping crane roosting than non-Program lands (dashed line).

What the science says in 2019:

- During the First Increment of the Program, tern and plover populations on the central Platte River have increased significantly (Pearson’s Rank Correlation test $\alpha \leq 0.05$) and proportionally to increases in off-channel nesting habitat creation efforts.
- On-channel nesting habitat creation efforts have not been successful and have largely been abandoned by terns and plovers.
- During the First Increment of the Program, UOCW on Program lands transitioned from significantly narrower than non-Program lands in 2010 to significantly wider in 2013 through 2019.
- There has been a discernable trend in the proportion of the whooping crane population roosting on Program lands since 2007. There has been a significant increase in the proportion of the population using the AHR as a whole during the spring migration through 2018, but not during the fall migration period.
- There has been a discernable trend in the proportion of the whooping crane population roosting within the AHR since 2007; however wet meadow use has remained steady and low.
- Whooping cranes have used off-channel palustrine wetland sites created and maintained by the Program during the First Increment; however wetland use has remained steady and low.

We estimate with confidence that:

- Program efforts to create and maintain off-channel tern and plover nesting habitat have been successful.
- Program efforts to create and maintain in-channel whooping crane roosting habitat have been successful.

Answering Big Question #10 during the First Increment:


- The Program has noted trends related to the tern and plover hypotheses in response to off-channel nesting habitat and has recently observed substantial use of Program and other conservation lands by whooping cranes.
- The Program is currently negotiating a 13-year Extension of the First Increment due in part to the fact that water objective of reducing annual shortages to target flows by 130,000-150,000 acre-feet has not been met.

Management implications:

- Synthesis of multiple lines of evidence related to this Big Question and the underlying system-level hypotheses have provided guidance to the GC regarding Program land and water management during the First Increment and will continue to into the Extension.

Priority Hypothesis	Results	
	LTPP	WC
S1a (habitat response)		
S1b (species response)		
S1c (wet meadows)		

Relationship to Big Question #10
It is hypothesized that Program management actions will result in measurable changes in central Platte River form and function, which will in turn result in detectable responses by the target species.



Additional First Increment Learning

Whooping Cranes

- Spring whooping crane migration season has shifted earlier in the year from 1975-1999 (95th percentile dates: March 20 – April 29) until 2007-2016 (current 95th percentile dates: March 6 – April 20).
- Fall whooping migration season has shifted later in the year from 1975-1999 (October 9 – November 10) until 2007-2016 (current 95th percentile dates: October 15 – November 16).
- Whooping crane migration corridor throughout the Great Plains is shifting eastward; however, use of the western portion of the AHR is increasing.
- No evidence of a relationship between flow and where whooping cranes choose to roost within the central Platte River.
- Whooping crane riverine habitat selection similar throughout the Great Plains and within the AHR.
- Whooping cranes select cornfields and riverine habitats over wet meadows within the AHR; however, throughout the Great Plains whooping cranes select riverine, open water, semi-permanent wetland, and wet meadow habitats over all upland landcover classes including cornfields.

Least Terns and Piping Plovers

- Least tern and piping plover monitoring from outside the nesting colony is sufficient for documenting abundance and productivity.
- The relative probability of use for least terns and piping plovers is maximized when distance to the nearest predator perch is ≥ 500 feet and elevation above the waterline is 7–10 feet.
- Probability of nesting by least terns increased as distance to water increased, whereas the probability of use by piping plovers was maximized when distance to water was ~ 150 feet.
- Piping plover territories are ~ 3 acres in size on off-channel sites.
- Productivity of least terns and piping plovers is reduced during both the nesting and brood rearing stages by weather factors more than factors the Program can manage.
- Electrified predator fences and wing panels are effective at deterring terrestrial predators from off-channel nesting areas.
- Habitat management activities implemented at off-channel sites to date seem to be sufficient for maintaining high levels of productivity for least terns and piping plovers within the AHR.
- The timing the spring pulse flow coincides with the least terns and piping plovers nesting season which limits the opportunity for successful nesting within the AHR.
- Least terns and piping plovers select channels that are $\geq 1,200$ feet wide in other river systems.

- Mechanical in-channel tern and plover nesting islands are difficult to construct, expensive, erode quickly during natural high flow events. Tern and plover productivity is also much lower than on off-channel sand and water (OCSW) sites.¹⁷
- The GC decided to forego any further in-channel island construction activities in lieu of creating an additional 60 acres of bare-sand OCSW nesting habitat and maintaining 10 acres of moving complex approach (MCA) nesting islands in the channel.
- In 2016, the Service concluded that creating an additional 60 acres of off-channel nesting habitat would meet their criterion of achieving “stable or increasing populations” of least terns and piping plovers within the AHR¹⁸.

Pallid Sturgeon

- No evidence pallid sturgeon use discharge as a spawning cue as the lower Platte River has been used during both low- and high-discharge years.
- All but one of the four gravid females known to have entered the Platte River to spawn did so when temperatures were $< 15^{\circ}\text{C}$ and exited when temperatures exceeded 20°C .
- No evidence that post-spawn females exit the lower Platte River due to low discharge as females generally exited the Platte River at a higher discharge than when they entered.
- Spawned pallid sturgeon exited the lower Platte River approximately one to two weeks after a peak flow event temporarily increased discharge and depressed water temperature.
- If one assumes diurnal water temperature variation is important for spawning behavior, the Program may be able to influence this metric through flow-releases to reduce variation. However, given the magnitude and duration of any potential flow release the Program could currently make, we found our potential to reduce diurnal variation in water temperature is minute in comparison the natural variability in diurnal water temperature variation and may only be detectable at the lowest flows observed in the lower Platte River.
- Most pallids were captured below the Elkhorn River at moderate discharges (6,000–10,000 cfs) with few pallids captured at very low or very high discharges. In addition, few pallids were captured upstream of the Elkhorn River, even in seasons with moderate discharges. The lack of any discernable relationship between discharge and captures makes it difficult to conclude that differences in discharge is the main driver of the disparity in captures above and below the Elkhorn River confluence.
- When capture effort is taken into account, several stretches of river above the Elkhorn river had similar captures rates as those below the Elkhorn River.
- The only habitat metric found to be markedly different upstream and downstream of the Elkhorn River confluence is turbidity.

Sandhill Cranes

- Sandhill cranes select wide unobstructed channels with low flows (i.e., 900-ft unobstructed channel with $\leq 1,400$ cfs).
- Spring sandhill crane migration season is shifting earlier in the year.
- Sandhill crane roosting within the AHR is shifting eastward.

Wet Meadows

- Whooping crane use of the AHR has increased during the spring migration while wet meadow use has remained stable and low.

System

- Remote channel measures are adequate for capturing channel width metrics such as maximum unvegetated channel width, total unvegetated channel width, total channel width, etc.












APPENDIX A



2019 State of the Platte Priority Hypotheses Status Table



2019 State of the Platte Priority Hypotheses Status Table. Status of AMP priority hypotheses, as listed in Table 2 of the Adaptive Management Plan (Page 70). See shape coding key at end of table.

X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
System							
S1	The Platte River form can be modified by either mechanical/sediment/flow management (i.e., clear/level/pulse) or mechanical means along with non-Program managed flows (i.e., clear/level/mechanical).		#10	Geomorphology and vegetation monitoring, LiDAR and other aerial imagery, EDO analyses		Collecting the data necessary to answer all S1 hypotheses. To date, <i>State of the Platte</i> evaluations focused on BQ #1-#10.	
S1a	Program channel habitat restoration actions will result in detectable change to Platte River form and function.	Cannot detect a significant effect on indicators.	#10	Geomorphology and vegetation monitoring, LiDAR and other aerial imagery, EDO analyses		<i>2016 State of the Platte</i> – During the First Increment of the Program, UOCW on Program lands transitioned from significantly narrower than non-Program lands in 2010 to significantly wider in 2013 through 2016.	
S1b	Program land management actions (i.e., restoration into habitat complexes) will have a detectable effect on target bird species use of the associated habitats.	Cannot detect a significant effect on indicators	#10	Geomorphology and vegetation monitoring, LiDAR and other aerial imagery, bird monitoring, EDO analyses		<i>2016 State of the Platte</i> – Monitoring and analyses indicate there is a strong positive correlation between Program-defined suitable nesting habitat and tern and plover breeding pair counts within the AHR. Less of a correlation for whooping cranes.	
S1c	Program actions will increase functional wet meadows in habitat complexes during the First Increment.		#10	N/A		<i>2016 State of the Platte</i> – TBD	
S2	Implementing Program land and water management actions (i.e., habitat complexes and clear/level/pulse) will have a detectable effect on other species use of the associated habitats.	Within the overall management objectives for whooping cranes, terns and plovers, and pallids sturgeon, benefits can be provided to non-target listed species and non-listed species of concern thereby reducing the likelihood of future listing and improve overall ecosystem diversity.	N/A	N/A			


X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Terns and Plovers							
T1	Additional bare sand habitat will increase the number of adult least terns.	Bare sand is not currently limiting number of adults.	#6	PRRIP tern/plover monitoring protocol, EDO analyses, tern/plover habitat synthesis chapters		<i>2015 State of the Platte</i> – Monitoring and analyses indicate there is a <u>strong positive correlation</u> between Program-defined suitable nesting habitat and tern and plover breeding pair counts within the AHR.	
T2	Tern productivity is related to the number of prey fish (<3 inches) and fish numbers limit tern production below 800 cfs from May-Sept.	Prey fish do not limit tern production at 799 cfs or tern production is limited by summer flows of <50 cfs.	#8	Districts' forage fish monitoring protocol, USGS foraging habits study, EDO analyses and publication		<i>2016 State of the Platte</i> – Monitoring and analyses indicate there is no relationship between tern use and productivity and flow (i.e. forage fish) within the AHR.	
T2a	Flow rates influence the number and species diversity in tern prey base (fish).	Tern productivity not affected by fish community species diversity.	N/A	N/A			
P1	Additional bare sand habitat will increase the number of adult piping plovers.	Bare sand is not currently limiting number of adults.	#6	PRRIP tern/plover monitoring protocol, EDO analyses, tern/plover habitat synthesis chapters and associated publications		<i>2015 State of the Platte</i> – Monitoring and analyses indicate there is a <u>strong positive correlation</u> between Program-defined suitable nesting habitat and tern and plover breeding pair counts within the AHR.	
P2	Plover productivity is related to the number of suitable macroinverts and macroinverts limit plover production below 800 cfs from May-Sept.	Macroinverts do not limit plover production at 799 cfs or plover production is limited by summer flows of <50 cfs.	#8	Districts' forage fish monitoring protocol, USGS foraging habits study, EDO analyses		<i>2016 State of the Platte</i> – Monitoring and analyses indicate there is likely no relationship between plover use and productivity and flow (i.e. forage fish) within the AHR.	



X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
TP 1	Interaction of river and sandpit habitat.	LT and PP show no preference for the river over sandpits.	#7	PRRIP tern/plover monitoring protocol, EDO analyses		<i>2015 State of the Platte</i> – Monitoring and analyses indicate both in-channel and off-channel nesting habitats are <u>not necessary</u> to maintain the central Platte River population of terns and plovers. However, the river is a valuable source of forage for both species as forage availability is lower on off-channel habitats.	
TP 2	The central Platte River may act as a source or sink for terns and plovers.	Currently not a sink.	N/A	PRRIP tern/plover monitoring protocol, EDO analyses		<i>2015 State of the Platte</i> – Given population growth within the AHR and fledge ratios that exceed all numbers hypothesized to result in population growth, the hypothesis is <u>rejected</u> .	
TP 4d	Correlation between river island habitat and flow.		N/A	Tern/plover habitat synthesis chapters		<u>No need to test</u> as sandbars are not suitably high for nesting.	
TP 5	Use of riverine islands by least terns and piping plovers will increase with active channel width.	Use will not increase with channel width.	#1	Tern/plover habitat synthesis chapters		<u>Hypothesis affirmed</u> in Tern/plover synthesis chapter 4.	


X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Whooping Cranes							
WC 1	Whooping crane use will increase as function of Program land and water management activities.	Whooping crane use will not increase as function of Program land and water management activities.	N/A	WEST habitat selection report, whooping crane habitat synthesis chapters		Hypothesis affirmed in 2017 and 2018 spring and fall whooping crane monitoring reports and several Program publications.	
WC 3	Whooping crane use is related to habitat suitability. The prediction of habitat suitability for whooping crane in channel habitat as a function of water depth (preferred depth?) and channel width (define as wetted width, open width, other?).	WC use of areas is not directly linked to FWS habitat suitability values.	#5	WEST habitat selection report, whooping crane habitat synthesis chapters		<i>2016 State of the Platte – Whooping cranes select channel widths of 650 ft and unforested corridor widths of 1,100 ft.</i>	
WC 4	Whooping crane use of the central Platte River study area will increase proportionally to an increase in wet meadows.	WC do not use wet meadows currently and are unlikely to respond to increases in wet meadow area.	N/A	N/A		Evidence points to <u>rejecting</u> this hypothesis. Peer review of key documents will likely result in a conclusive answer in a future <i>State of the Platte Report</i> .	
WC 5	Whooping cranes are adversely affected by nocturnal disturbances that lead to flushing (walking or flying) which could lead to potential mortality.	WC are not negatively impacted by nocturnal disturbances.	N/A	N/A			



X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Pallid Sturgeon							
PS-1	Program flow/sediment management will result in a positive species response by the pallid sturgeon in the lower Platte River.	Program flow/sediment management will result in no increase in species use/occurrence by the plaid sturgeon in the lower Platte River.	N/A	N/A		The Program is in the process of determining next steps on pallid sturgeon Big Questions, hypotheses, and issues for the First Increment and First Increment Extension through the facilitated Pallid Sturgeon Process. Determining linkages between Program hypotheses, management actions, and pallid sturgeon response will be part of that process.	
PS-2	Program water management will result in measurable changes on flow in the lower Platte River.	Program water management will result in statistically insignificant changes on flow in the lower Platte River.	#9	Stage change study		The Program is in the process of determining next steps on pallid sturgeon Big Questions, hypotheses, and issues for the First Increment and First Increment Extension through the facilitated Pallid Sturgeon Process. Determining linkages between Program hypotheses, management actions, and pallid sturgeon response will be part of that process.	
PS-4	Flows in the lower Platte will affect pallid sturgeon habitat suitability.	Flows in the lower Platte River will have no effect on pallid sturgeon habitat suitability.	N/A	N/A		The Program is in the process of determining next steps on pallid sturgeon Big Questions, hypotheses, and issues for the First Increment and First Increment Extension through the facilitated Pallid Sturgeon Process. Determining linkages between Program hypotheses, management actions, and pallid sturgeon response will be part of that process.	




X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
PS-5	Pallid sturgeon habitat suitability is maximized between water temperatures of X and Y in the lower Platte River.	Pallid sturgeon use is independent of river water temperature.	N/A	N/A		The Program is in the process of determining next steps on pallid sturgeon Big Questions, hypotheses, and issues for the First Increment and First Increment Extension through the facilitated Pallid Sturgeon Process. Determining linkages between Program hypotheses, management actions, and pallid sturgeon response will be part of that process.	
PS-6	Increasing flow in the lower Platte will affect pallid sturgeon habitat availability.	Increasing flow in the lower Platte River will have no effect on pallid sturgeon habitat availability.	N/A	N/A		The Program is in the process of determining next steps on pallid sturgeon Big Questions, hypotheses, and issues for the First Increment and First Increment Extension through the facilitated Pallid Sturgeon Process. Determining linkages between Program hypotheses, management actions, and pallid sturgeon response will be part of that process.	
PS-7	Increasing habitat availability in the lower Platte will increase pallid sturgeon use.	Pallid sturgeon use is independent of lower Platte River habitat availability.	N/A	N/A		The Program is in the process of determining next steps on pallid sturgeon Big Questions, hypotheses, and issues for the First Increment and First Increment Extension through the facilitated Pallid Sturgeon Process. Determining linkages between Program hypotheses, management actions, and pallid sturgeon response will be part of that process.	





X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
PS-9	Increasing Program flow releases will decrease water temperatures in the lower Platte River.	River water temperature is independent of flow rate in the lower Platte River Increases in program flow releases will increase water temperatures on the lower Platte River.	N/A	N/A			
PS-11	Non-Program actions (e.g. harvest, stocking, Missouri River conditions) determine the occurrence of pallid sturgeon in the lower Platte River.	Program actions will affect the rate of occurrence of pallid sturgeon in the lower Platte River such that use is disproportionate to external factors (e.g., stocking, harvest, local conditions) relative to local population.	N/A	N/A			
Physical Processes – Flow							
Flow #1	Increasing the variation between river stage at peak (indexed by Q1.5 flow at Overton) and average flows (1,200 cfs index flow), by increasing the stage of the peak (1.5-yr) flow through Program flows, will increase the height of sand bars between Overton and Chapman by 30% to 50% from existing conditions.	Flow magnitudes and channel compilations are insufficient to generate bars high enough to provide habitat for LT and PP. Bars may quickly vegetate making them poor habitat for target species. Bars can be created/maintained by mechanical/other means.	#1	Geomorphology and vegetation monitoring, tern/plover monitoring, tern/plover habitat synthesis chapters		<i>2016 State of the Platte</i> – Full SDHF magnitude of 8,000 cfs is not sufficient to create sandbars exceeding the PRRIP’s minimum height suitability criterion. Sandbars created by SDHF releases will be inundated during the nesting season in most years.	

X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Flow #3	Increasing 1.5-yr Q with Program flows will increase local boundary shear stress and frequency of inundation at existing green line (elevation at which riparian vegetation can establish). These changes will increase riparian plant mortality along margins of channel, raising elevation of green line. Raised green line = more exposed sandbar area and wider unvegetated main channel.	Insufficient Program flows to adequately increase shear stress on banks. Plant mortality can be achieved by other means.	#2	Directed scour research, whooping crane habitat synthesis chapters		<i>2016 State of the Platte</i> – The comparatively short duration and low volume of SDHF limits the predicted increase in UOCW to ≤12 ft. SDHF duration is not sufficient to create and maintain UOCWs that are suitable for whooping crane roosting.	
Flow #4	Annual riparian seedling mortality greater than 90% is required to prevent riparian encroachment on exposed bars, thereby increasing (maintaining at least 10 acres/mile) exposed bars between Overton and Grand Island that are usable as LT and PP habitat.	Riparian seedling mortality greater than 90% is needed to increase exposed bar area. Other factors drive exposed bar area instead of seedling mortality. Plant mortality can be achieved by other means.	N/A	N/A			
Flow #5	Increasing magnitude and duration of a 1.5-yr flow will increase riparian plant mortality along the margins of the river. There will be different relations (graphs) for different species.	Insufficient Program flows to maintain required flow durations. Plant mortality can be achieved by other means.	#2	Directed scour research, whooping crane habitat synthesis chapters		<i>2016 State of the Platte</i> – Mature phragmites plants or plant patches have a very low probability of being eroded at the highest flow magnitudes and velocities observed in the AHR. An herbicide control program is ongoing	

X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Physical Processes – Sediment							
Sediment #1	Average sediment augmentation at Overton of 185,000 tons/yr. under existing flow regime and 225,000 tons/yr. under Governance Committee proposed flow regime achieves a sediment balance to Kearney.	Augmentation greater than or less than 225,000 tons/year is needed to balance the sediment budget and increase exposed bar area. There is no sediment imbalance. Exposed bar area or occurrence of braiding will not be affected by increased sediment. Sediment balance is insignificant except in local instances. Satisfactory bar areas can be created and maintained through strictly mechanical actions.	#3	Sediment transport modeling, results of sediment augmentation Proof of Concept experimental implementation		Augmentation of sediment in the south channel is necessary to slow incision and narrowing and prevent degradation from progressing downstream past the Overton bridge. It will be challenging to measure the effectiveness of augmentation given that the desired beneficial effect is slowing and ultimately halting of a long-term trend.	
Sediment #2	A balanced sediment budget (sediment augmentation of 225,000 tons/year near Overton under proposed Governance Committee flows) when implemented with mechanical actions (channel consolidation & widening) in anastomosed reaches will promote braided channel morphology with an average braiding index in the main channel of greater than 3.	Flows and sediment augmentation are insufficient to achieve desired braiding index.	N/A	N/A			

X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Sediment #3	Increasing the average braiding index of the main channel by achieving a balanced sediment budget, increases the active unvegetated width of the main channel at an index flow of 2,000 cfs (at Overton).	Width will not change with increasing braiding index.	N/A	N/A			
Sediment #4	Increasing the average braiding index to greater than 3 for the main channel in the sediment deficient reach near Overton will increase and maintain exposed bar area greater than 1.5 acres in the reach between Overton and Kearney at an index flow of 1,200 cfs (at Overton).	There is no relationship between braiding index and area of exposed bars. Exposed bars may be created (maintained) through mechanical means without need to change braiding index.	N/A	N/A			
Physical Processes – Mechanical							
Mechanical #2	Increasing the Q1.5 in the main channel by consolidating 85% of the flow, and aided by Program flow and a sediment balance, flows will exceed stream power thresholds that will convert main channel from meander morphology in anastomosed reaches, to braided morphology with an average braiding index > 3.	Higher stream power (higher 1.5 yr. Q and/or more consolidation of side channels) needed to convert channel to braided morphology. Lower stream power will convert channel to braided morphology.	#4	Directed scour research, whooping crane habitat synthesis chapters		2016 State of the Platte – Peak flows in the AHR are generally not sufficient to remove mature woody vegetation or erosion-resistant species like phragmites. Mechanical clearing and leveling are necessary to create suitable channel configurations and facilitate channel adjustments to changes in flow and sediment.	

X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Mechanical #3	Reducing the number of channels in a transect to 3 or less under balanced sediment budget will convert anastomosed reaches of the Platte River between Overton and Chapman to a braided channel morphology. With proposed flow regime, should occur with greater number of channels.	Reducing the number of channels in a transect to 1 or 2 is necessary to achieve an average braiding index in the main channel of greater than 3.	N/A	N/A			
Mechanical #4	Increasing the average braiding index to greater than 3 in the main channel by channel manipulation will promote in the Platte River at the mechanically changed sites a total main channel wetted width exceeding 500 to 750 ft at an index flow of 1,700 cfs (at Overton).	A braiding index greater than 4 is needed to achieve a width greater than 500 ft. There is no relation between braiding index and channel width.	N/A	N/A			
Mechanical #5	Increasing the average braiding index to greater than 3 for the main channel by mechanical channel manipulation, will increase and maintain exposed bar area greater than 1.5 acres at mechanical changed sites at an index flow of 1,200 cfs (at Overton).	Mechanically consolidating flows will have no effect on areal extent of bars.	N/A	N/A			

X-Y Graph Number	Description of hypothesis	Description of alternative/competing hypotheses	Link to PRRIP Big Questions	Data Source(s)	Conclusion ⁱ	Notes	Comments from TAC/AMWG
Wet Meadows							
WM-2	Wet meadows producing the optimum productivity and diversity of macro-invertebrates potentially consumed by WC exhibit certain characteristic combinations of soils, hydrology, size and location. Mormon Island and adjacent to Rowe Sanctuary have some of best existing combinations	There are too many possible combinations of site characteristics to allow for a meaningful characterization of “desirable” conditions.	N/A	N/A			
WM-3	Shallow surface water and groundwater in March and April support high productivity and diversity of macroinvertebrates as potential food sources to WC in wet meadows.		N/A	N/A			
WM-4	A predominance of organic-rich soils supports the productivity and diversity of macro-invertebrates as potential WC food sources in bottomland grasslands.	Wet meadows and their soils are too complex and variable to allow this individual factor to be effectively assessed.	N/A	N/A			
WM-8a	As the spring depth to groundwater increases, surface soils stay frozen longer. Where groundwater is closer to the surface soils thaw sooner.		N/A	N/A			

ⁱ Conclusions are indicated as one of the following categories:



Hypothesis answered conclusively – affirmed.



Hypothesis answered conclusively – rejected.



Hypothesis not yet answered – ongoing implementation, analysis, and synthesis.



Not currently being addressed through implementation of the AMP and related data analysis and synthesis.



APPENDIX B

PRRIP Peer Reviewed Papers & Reports

REFEREED PUBLICATIONS

Baasch, D.M., P.D. Farrell, A.J. Caven, K.C. King, J.M. Farnsworth, and C.B. Smith. 2019. [Sandhill crane use of riverine stopover sites along the central Platte River in Nebraska, USA](#). Monographs of the Western North American Naturalist 11:1–13.

Baasch, D.M., P.D. Farrell, A.T. Pearse, D.A. Brandt, A.J. Caven, M.J. Harner, G.D. Wright, and K.L. Metzger. 2019. [Diurnal habitat selection of migrating whooping cranes in the Great Plains](#). Avian Conservation and Ecology. 10.5751/ACE-01317-140106.

Baasch, D.M., P.D. Farrell, S. Howlin, A.T. Pearse, J.M. Farnsworth, and C.B. Smith. 2019. [Whooping Crane use of riverine stopover sites](#). PLOS ONE 14 (1): e0209612. <https://doi.org/10.1371/journal.pone.0209612>.

Farnsworth, J.M., D.M. Baasch, P.D. Farrell, and C.B. Smith. 2018. [Investigating whooping crane habitat in relation to hydrology, channel morphology and a water-centric management strategy on the central Platte River, Nebraska](#). Heliyon. <https://doi.org/10.1016/j.heliyon.2018.e00851>.

Farnsworth, J.M., D.M. Baasch, and P.D. Farrell. 2018. [Integrating species-centric and geomorphic-centric views of interior least tern and piping plover habitat selection](#). Heliyon. <https://doi.org/10.1016/j.heliyon.2018.e00648>.

Farnsworth, J.M., D.M. Baasch, C.B. Smith, and K.L. Werbylo. 2018. [Reproductive ecology of interior least tern and piping plover in relation to Platte River hydrology and sandbar dynamics: Response to the Letter to the Editor](#). Ecology and Evolution. <https://doi.org/10.1002/ece3.4097>

Farrell P.D., D.M. Baasch, J.M. Farnsworth, and C.B. Smith. 2018. [Least tern and piping plover nest success and brood survival at off channel sites in the Central Platte River](#). Avian Conservation and Ecology 13:1. <https://doi.org/10.5751/ACE-01133-130101>.

Baasch, D.M., P.D. Farrell, J.M. Farnsworth, and C.B. Smith. 2017. [Nest-site selection by Interior Least Terns and Piping Plovers at managed, off-channel sites along the Central Platte River in Nebraska, USA](#). Journal of Field Ornithology 88:236–249.

Farnsworth, J.M., D.M. Baasch, C.B. Smith, and K.L. Werbylo. 2017. [Reproductive ecology of interior least tern and piping plover in relation to Platte River hydrology and sandbar dynamics](#). Ecology and Evolution 10:3579–3589. doi: 10.1002/ece3.2964.

Baasch, D.M., P.D. Farrell, J.M. Farnsworth, and C.B. Smith. 2017. [Interior least tern productivity in relation to flow in the central Platte River valley](#). Great Plains Research 27:35–42.

Smith, C.B., J.M. Farnsworth, D.M. Baasch, and J.F. Kenny. 2016. [Adaptive Management and Governance Lessons from a Semiarid River Basin: A Platte River Case Study](#). In: Miller, K.A., Hamlet, A.F., Kenney, D.S., Redmond, K.T. (eds.), Water Policy and Planning in a Variable and Changing Climate: Insights from the Western United States, CRC Press.

Pearse, A.T., M.J. Harner, D.M. Baasch, G.D. Wright, A.J. Caven, and K.L. Metzger. 2016. [Evaluation of nocturnal roost and diurnal sites used by whooping cranes in the Great Plains, United States](#). Open File Report 2016–1209.

Werbylo, K.L., J.M. Farnsworth, D.M. Baasch, and P.D. Farrell. 2016. [Investigating the accuracy of estimated unvegetated channel widths in a braided river system: a Platte River case study](#). Geomorphology 278:163–178. <https://doi.org/10.1016/j.geomorph.2016.11.003>

Farnsworth, J.M., J.F. Kenny, and C.B. Smith. 2015. [Comment on “Progressive abandonment and planform changes of the central Platte River in Nebraska, central USA, over historical timeframes”](#). Geomorphology 250:437–439. <https://doi.org/10.1016/j.geomorph.2014.12.014>

Pearse, A.T., D.A. Brandt, W.C. Harrell, K.L. Metzger, D.M. Baasch, and Hefley, T.J., 2015, [Whooping crane stopover site use intensity within the Great Plains: U.S. Geological Survey Open-File Report 2015–1166](#), 12 p., <http://dx.doi.org/10.3133/ofr20151166>.

Baasch, D.M., T.J. Hefley, and S.D. Cahis. 2015. [A comparison of breeding population estimators using nest and brood monitoring data](#). Ecology and Evolution 5:4197–4209. doi: 10.1002/ece3.1680

Hefley, T.J., D.M. Baasch, A.J. Tyre and E.E. Blankenship. 2015. [Use of opportunistic sightings and expert knowledge to predict and compare Whooping Crane stopover habitat](#). Conservation Biology 29:1337–1346. doi: 10.1111/cobi.12515

Hefley, T.J., D.M. Baasch, A.J. Tyre, and E.E. Blankenship. 2014. [Correction of location errors for species distribution models](#). Methods in Ecology and Evolution 5:207–214.

Hefley, T.J., A.J. Tyre, D.M. Baasch and E.E. Blankenship. 2013. [Non-detection sampling bias in marked presence-only data](#). Ecology and Evolution 3:5225–5236.

Smith, C.B. 2011. [Adaptive management on the central Platte River – science, engineering, and decision analysis to assist in the recovery of four species](#). Journal of Environmental Management 92:1414–1419.

Smith, C.B. 2009. [Active adaptive management on the Platte River](#). Water Resources IMPACT, Journal of the American Water Resources Association 11:8–10.

PEER REVIEWED SYNTHESIS REPORTS

Executive Director's Office. 2017. [Whooping Crane Habitat Synthesis Chapters](#). Prepared for the Governance Committee of the Platte River Recovery Implementation Program.

Executive Director's Office. 2015. [Interior Least Tern and Piping Plover Habitat Synthesis Chapters](#). Prepared for the Governance Committee of the Platte River Recovery Implementation Program.



APPENDIX C

PRRIP Progress - Land & Water Objectives



Land Objective

The Program’s First Increment land objective is to acquire, protect, and restore 10,000 acres of habitat for the three target avian species.

As of 2019, the Program has exceeded the 10,000-acre First Increment land objective by 2,783 acres. Most of the Program’s habitat lands are located in blocks referred to as habitat complexes. There are six primary habitat complexes (Cottonwood Ranch, Elm Creek, Pawnee, Fort Kearny, Clark Island, and Shoemaker Island) with additional complex habitat at four other locations (Plum Creek, Minden to Gibbon, Alda to Grand Island, and Grand Island to Chapman). In addition, the Program manages 645 acres of non-complex lands comprised of four OCSW and two palustrine wetland sites as well as 25 acres set aside for Clean Water Act compliance. A map of Program properties can be found on page 2 of this report.

Habitat Lands	Number of Acres
Plum Creek	866
Cottonwood Ranch	3,552
Elm Creek	1,438
Pawnee	742
Fort Kearny	2,190
Gibbon to Minden	834
Clark Island	784
Shoemaker Island	1,411
Alda to Grand Island	286
Grand Island to Chapman	10
Total Complex Land	12,113
Off-Channel Sand and Water	391
Palustrine Wetlands	254
Total Non-Complex Land	645
Clean Water Act Land	25
Grand Total	12,783

Water Objective

The Program’s First Increment water objective is to reduce deficits to USFWS target flows by an average of 130,000 - 150,000 acre-ft annually.

As of 2019, Program water projects reduce deficits to USFWS target flows by an average of over 112,000 acre-ft annually. This includes the three original state projects with a total score of 80,000 acre-feet per year (AFY) and the five-scored water action plan (WAP) projects (Phelps County Canal Recharge, Cook Recapture Well, Pathfinder Municipal Lease, No-Cost Net Controllable Conserved, and Elwood Recharge) with a score of 12,370 AFY. Other WAP projects are active but have yet to be scored including Central Nebraska Public Power and Irrigation District surface water and recharge, Nebraska Public Power District recharge, and CPNRD Surface Water Exchange. It is anticipated that these projects will increase the water project score by approximately 20,000 AFY. A large portion of the remaining First Increment water objective was going to be provided by the J-2 Regulating Reservoir, but this project has been put on hold due to cost, land acquisition and other concerns. Consequently, the water objective will not be met prior to 2019 and Program Signatories have agreed to pursue a 13-year First Increment Extension. Projects to be implemented during the First Increment Extension include but are not limited to: broad-scale recharge projects, slurry wall storage facilities, and acquire and retire. These projects will be used to achieve a total score of 120,000 AFY. Research will then be conducted to evaluate the need for the remaining 10,000 acres prior to implementation of the projects that would be necessary to achieve the First Increment water objective.

Water Project	Score (AFY)
Three State Projects	80,000
Phelps Co. Canal Recharge	2,700
Cook Recapture Well	160
Pathfinder Municipal Lease	6,350
No-Cost Net Controllable Conserved	260
Elwood Recharge	2,800
CNPPID Irrigator Lease	1,900
Total	94,170
Other operational WAP projects that have yet to be scored	~20,000

ENDNOTES

- ¹ Program. 2016. Analysis of sandbar height distributions following First Increment peak flow events. Prepared for the Platte River Recovery Implementation Program.
- ² Program. 2015. Least tern and piping plover habitat synthesis chapters. Prepared for the Platte River Recovery Implementation Program.
- ³ Bankhead et al. 2016. A combined field, laboratory and numerical study of the forces applied to, and the potential for removal of, bar top vegetation in a braided river. *Earth Surface Processes and Landforms*, 42(3), 439-459.
- ⁴ Tetra Tech Inc. 2017. 2016 Final Data Analysis Report Platte River Geomorphology and In-Channel Vegetation. Prepared for the Platte River Recovery Implementation Program
- ⁵ Program. 2017. Whooping Crane Data Synthesis Chapters. Prepared for the Platte River Recovery Implementation Program.
- ⁶ Tetra Tech Inc. 2015. Model Results, Platte River Sediment-transport Modeling, South Channel at Jeffery Island. Prepared for the Platte River Recovery Implementation Program.
- ⁷ Program. 2016. Proof-of-Concept Sediment Augmentation Implementation and Effectiveness Analyses Methods. Prepared for the Platte River Recovery Implementation Program.
- ⁸ Pollen-Bankhead et al. 2012. Can Short Duration High Flows be used to Remove Vegetation from Bars in the Central Platte River. Prepared for the Platte River Recovery Implementation Program.
- ⁹ See WEST Report titled Correlates of Whooping Crane Habitat Selection and Trends in Use in the Central Platte River 02-16-16.
- ¹⁰ See the Whooping Crane Synthesis Chapters 2 and 3.
- ¹¹ Baasch et al. 2019. Whooping Crane use of riverine stopover sites
- ¹² See PRRIP 2016 Tern and Plover Monitoring Report.
- ¹³ Baasch et al. 2015. A comparison of breeding population estimators using nest and brood monitoring data.
- ¹⁴ Baasch et al. 2017. Nest-site selection by Interior Least Terns and Piping Plovers at managed, off-channel sites along the Central Platte River in Nebraska, USA.
- ¹⁵ Sherfy et al. 2012. Foraging Ecology of Least Terns and Piping Plovers Nesting on Central Platte River Sandpits and Sandbars.
- ¹⁶ Baasch et al. 2017. Interior Least Tern Productivity in Relation to Flow in the Central Platte River.
- ¹⁷ Final PRRIP 2016 Tern and Plover SDM Report
- ¹⁸ Final PRRIP 2016 Tern and Plover SDM Report